



A.D. 1308
unipg

DIPARTIMENTO
DI SCIENZE AGRARIE,
ALIMENTARI E AMBIENTALI

An index for adaptability evaluation of slow-growing chicken genotypes reared in free-range system

S. Mattioli, L. Menchetti, E. Angelucci, A. Cartoni Mancinelli, A. Dal Bosco, L. Madeo, D. Chiattelli, F. Di Federico, C. Castellini




75th EAAP Annual Meeting
1/5 September 2024 - Florence, Italy




Outdoor adaptability criteria

Extensive Rearing System (ERS) must **optimize a production system that promotes biodiversity, environmental sustainability and food safety** (*National Organic Standards Board, 1995*)



What means suitable broiler genotypes adapted to outdoor system?

ADAPTABILITY




- Mancinelli, A. C., Amato, M. G., Zillio, D. M., Dal Bosco, A., Mattioli, S., & Castellini, C. (2017). Chicken adaptability in alternative systems adaptive response of chicken strains to the organic and free range rearing systems. *J. Dairy Vet. Sci*, 4(4).
- Guarino Amato, M., & Castellini, C. (2022). Adaptability challenges for organic broiler chickens: a commentary. *Animals*, 12(11), 1354.
- Cartoni Mancinelli, A., Mattioli, S., Menchetti, L., Dal Bosco, A., Ciarelli, C., Guarino Amato, M., & Castellini, C. (2021). The assessment of a multifactorial score for the adaptability evaluation of six poultry genotypes to the organic system. *Animals*, 11(10), 2992.
- Castellini, C., Mugnai, C., Moscati, L., Mattioli, S., Guarino Amato, M., Cartoni Mancinelli, A., & Dal Bosco, A. (2016). Adaptation to organic rearing system of eight different chicken genotypes: behaviour, welfare and performance. *Italian journal of animal science*, 15(1), 37-46.
- Cartoni Mancinelli, A., Mattioli, S., Dal Bosco, A., Aliberti, A., Guarino Amato, M., & Castellini, C. (2020). Performance, behavior, and welfare status of six different organically reared poultry genotypes. *Animals*, 10(4), 550.
- Mattioli, S., Mancinelli, A. C., Menchetti, L., Dal Bosco, A., Madeo, L., Amato, M. G., ... & Castellini, C. (2021). How the kinetic behavior of organic chickens affects productive performance and blood and meat oxidative status: A study of six poultry genotypes. *Poultry Science*, 100(9), 101297.

MATERIAL & METHODS



RedJ (RJ)



Naked neck (NN)



Lohmann Dual (LD)



Robusta Maculata x
Sassò (CB)

- 100 hundred chicks/strain of both sexes were used.
- Chickens were reared in pens (2 pens/strain) with an indoor (0.10 m²/bird) and outdoor (4 m²/bird) area.
- The animals fed *ad libitum* the same starter and grower diets.
- At 81 days of age, 15 chickens/pen were selected and slaughtered.
- Many variables have been evaluated:
 - *On-farm* (productive performances, behaviors, lesions and feather conditions)
 - *In vivo* parameters (blood)
 - Carcasses and Meat cuts (breast and drumstick)

MATERIAL & METHODS: pillars

Behaviour	Performance	Health status	Physical -chemical traits	Meat oxidative status	Meat fatty acids profile and indexes	Blood fatty acids profile	Blood traits
Running	Carcass Weight	Plantar Lesions	Drip loss	Lipids (drumstick)	PUFA (breast)	PUFA blood	HCT (%)
Attack	Drumstick (no bone)	Sternal Lesions	(drumstick)	MDA (breast)	C18:2n-6, LA (breast)	SFA blood	Lysozyme
Swell	Breast Weight	Neck score	b* colour (breast)	Retinol (breast)	C18 (breast)	C18 blood	ROMS
Rest	Bust Yield	Breast score	Drip loss (breast)	Retinol (drumstick)	C18:1n-9 (breast)	n-6 blood	PAO
Allo-grooming	Breast Yield	Wings score	WHC (breast)	Tocols (drumstick)	MUFA (breast)	C18:2cis n-6, LA blood	Heterophiles/lymphocytes
Grooming	Bone Weight	Beck score	b* colour	Tocols (breast)	C20:2 (breast)	C16 blood	
Escape	Live Weight		(drumstick)	Total lipids (breast)	n-3 (breast)	C20:4n-6, AA blood	
Grass Pecking	Tibia Length		White Striping (WS)	Carbonyls (drumst.)	C20:5n-3, EPA (breast)	C20:5n-3, EPA blood	
Walking	Breast Thickness		a* colour	Carbonyls (breast)	C22:2 (breast)	C18:3 n-3, α-ALA blood	
Stretching	Sternum Length		(drumstick)		C22:5n-3, DPA (breast)	n-3 blood	
Hide			L* (drumstick)		C22:4 (breast)	C18:1n-9 blood	
Other_peck.			a* colour (breast)		C20:4n-6, AA (breast)	MUFA	
Sand					C14 (breast)	MDA blood	
Scratch					C18:3n-3, α-ALA (breast)		
					C16:1 (breast)		
					C17:1 (breast)		
					C14 (drumstick)		
					C22:6n-3, DHA (breast)		
					C14:1 (drumstick)		
					C16 (drumstick)		
					C16:1 (drumstick)		
					C20:5n-3, EPA (drumstick)		
					C18 (drumstick)		
					C22:2 (drumstick)		
					n-3 (drumstick)		
					PUFA (drumstick)		
					C22:6n-3, DHA (drumstick)		
					C18:1cis9 n-9 (drumstick)		
					C22:4 (drumstick)		
					C22:5n-3, DP (drumstick)		
					C20:4n-6, AA (drumstick)		

Simona Mattioli

75th EAAP Annual Meeting, 1/5 September 2024 - Florence, Italy

Discriminant Analysis

- Selection of the variables characterizing the genotype for 4 pillars

Principal Component and Reliability Analyses

- Creation of a composite Index and its validation and refinement


Graph

- Index scores and differences among genotypes

Cronbach alpha

- Refine the Index and assess its internal consistency reliability.

1. DA: selection of the variables characterizing the genotype for each pillar




- ✓ *stepwise method* (F value = 0.02)
- ✓ To avoid multicollinearity, the variables were first selected using correlation and pooled within-groups correlation matrices by eliminating those with a coefficient $\geq |0.8|$
- ✓ In the "behavior" pillar, variables with a mean occurrence < 1 were not included (i.e., rare behaviors).

- The DA produced the discriminant function -*most parsimonious linear combinations of indicators describing between-genotype differences for each pillar*-
- For each pillar, the variables to be included in subsequent analyzes were selected based on their contribution to discrimination (Dfs explaining at least 80% of the variance between genotypes).

2. Principal Component and Reliability Analyses: creation of a composite Index and its validation and refinement

DA _{behaviour}		DA _{body condition}		DA _{meat quality}			DA _{In vivo health}		
Variable	Df1 (98.8%)	Variable	Df1 (84.0%)	Variable	Df1 (55.5%)	Df2 (27.8%)	Variable	Df1 (65.6%)	Df2 (28.1%)
Resting	5.699	Live weight	0.724	HF12_breast	0.479	0.523	HCT (%)	1.101	-0.981
Scratching	4.066	DFI	0.457	Carbonyls_drumstick	0.460	0.225	tocols	0.603	0.053
Roosting	0.698	Breast yield	0.303	DRIP loss %_drumstick	0.319	-0.493	Retinol	0.488	0.156
Grass peking	0.114	Back Score	0.203	Carbonyls_breast	0.299	0.246	H/L	-0.057	0.547
Attacking	-3.712	Plantar Score	0.044	MUFA_breast	0.295	0.237	Carbonyls	-0.124	0.526
Grooming	-4.450	Breast Score	-0.470	pH_breast	0.268	-0.261	Lisozima	-0.129	-0.447
				n-3_drumistick	0.167	0.384	HGB	-1.266	0.351
				SFA_drumistick	-0.045	-0.614			
				n-6_drumistick	-0.083	-0.703			
				SFA_breast	-0.237	0.434			
				DRIP loss%_breast	-0.257	0.043			
				Retinol_breast	-0.352	-0.415			
				lipids_breast	-0.594	0.578			


RESULTS



Genotype_cod

- CB
- LD
- NN
- RJ
- Group Centroid


DA_{behaviour}



Genotype_cod

- CB
- LD
- NN
- RJ
- Group Centroid


DA_{meat quality}



Genotype_cod

- CB
- LD
- NN
- RJ
- Group Centroid

DA_{body condition}



Genotype_cod

- CB
- LD
- NN
- RJ
- Group Centroid

DA_{In vivo health}

3. Principal Component and Reliability Analyses: creation of a composite Index and its validation and refinement

REFINE

- One variable was excluded because it did not present any correlation coefficient >0.3 (SFA_drumstick).
- Other variables were removed because had a loading <0.32 (HGB, Scratching, Attack, Tocols_blood, n-6 and n-3 drumstick, retinol_breast, tocals_blood, lysozyme)


Component Matrix

	Component 1
Zscore (live weight)	.919
Zscore (DFI)	.876
Zscore (BREAST feather condition)	-.862
Zscore (lipids_breast)	.746
Zscore (breast yield)	.662
Zscore (DRIP loss %_drumstick)	-.655
Zscore (HFI2_breast)	-.535
Zscore (SFA_breast)	.521
Zscore (Confort)	-.499

FINAL SELECTION:

- 9 variables were thus selected for the creation of the Index.
- The Cronbach's Alpha value for this composite index was 0.869.
- A final PCA was conducted including the nine variables listed in Table (PCA explained 38.6% of the variance)

4. Index scores and differences among genotypes



CONCLUSIONS - Take home message

- ❖ A simple index for describing the chickens adaptability to ERS is needed
- ❖ The choice of the criteria for the index building is very important, more criteria produce more robust index. It is need
 1. Consistency: Pillars representability
 2. Reliability: number of variables to include
 3. Simplicity: few, simple and «cheap» variables to assess, with the aim to use them also in large-scale/on-farm application
- ❖ To reduce the criteria number is needed to refine the index by including more data (chicken genotypes) in the analysis

The tested genotypes are SG, therefore very similar in characteristics, consequently the variance explained by the criteria is limited.

Probably applying the index to extreme lines would be more explanatory.

PPILOW PARTNERS



Thank you for your attention

simona.mattioli@unipg.it

www.ppilow.eu